


1/28



Unmethylated CpG dinucleotide

K motif : contains essential sequences of TCGTA and TCGTT

activate monocytes / B cells, secrete IL-6

Important where antibody response is essential

D ODN: containing palindromic sequence such as ATCGAT
activating NK cells / secreting IFN- γ

longer sequence : e.g. GGTGCATCGATGCAGGGGGG

ODN sequence containing GACGTT shows optimal response in mouse

ODN sequence containing GTCGTT shows optimal response in human

FIG. 1 Structure-activity relationship of unmethylated CpG-containing nucleotide sequence

BEST AVAILABLE COPY

2/28




FIG. 2 CpG dinucleotide modified at 3'-end with various lipophilic group

3/28




FIG. 3 Modified CpG dinucleotide as di-valent ligand

4/28




FIG. 4 Hexa-nucleotide ATCGAT modified at 5'-end with a lipophilic group

5/28





FIG. 5 Glycerol Nucleic Acid (GNA) and peptide nucleic acid (PNA) as structural mimetics of DNA

6/28






FIG. 6 Hexa-nucleotide GTcgTT modified at 3'-end with a lipophilic group, wherein cg di-nucleotide has glycerol-based backbone.


8/28




dA-CE phosphoramidite




dT-CE phosphoramidite



dC-CE phosphoramidite



dG-CE phosphoramidite



Spacer-18 phosphoramidite

FIG. 8 Building blocks for solid-phase nucleotide synthesis by phosphoramidite method

9/28




FIG. 9 Modification of lcaa-CPG (long chain aminoalkyl controlled pore glass) resin for the synthesis of lipided oligonucleotides.

10/28




FIG. 10 Preparation of lipidated CpG dinucleotide 1 on solid phase

11/28





FIG. 11 Preparation of pentaerythritol-derived diolpo-alcohol 11 and its application for the synthesis of lipidated oligonucleotides (16)

12/28



- i. BnBr, NaH, DMF, RT, 94%
 ii. HOAc-H₂O, 40 °C, 85%
 iii. BzCN, Et₃N, CH₃CN, -50 °C, 60%
 iv. paraformaldehyde, DCM, HCl(g), 0 °C
 v. NaH, DMF, RT, 30%; or
 BSA, TBAI, DCM, RT, 35%
 vi. Me₂BBr, DCM, 0 °C, 76%; or
 BCl₃, DCM, -78 °C, 76%
 vii. (CNCH₂CH₂O)P'Pr₂-Cl, DIPEA, DCM, RT, 79%

FIG. 12 Preparation of glycerol-cytosine phosphoramidite 25

13/28





FIG. 13 Preparation of glycerol-guanosine building block 32

14/28

FIG. 14 Synthesis of glycerol-based CpG dinucleotide **8**

15/28




FIG. 15 Preparation of PNA-based CpG analogue 7 by standard solid phase peptide synthesis using Fmoc/Bhoc chemistry.

16/28





FIG. 16. Immunostimulatory adjuvant properties of CpG analogues 1-6. In vitro antigen specific proliferation of T cells from C57BL/6 mice immunized with a single dose of BLP25 liposomal vaccine formulation. The vaccine dose contains 20 μ g of MUC1-derived 25-mer lipopeptide as an antigen and 10 μ g of one of synthetic CpG analogues 1-6 as an adjuvant. R595 lipid A is used for comparison, which is the natural detoxified lipid A product isolated from *Salmonella minnesota* R595 and is currently being evaluated as a vaccine adjuvant in clinic.

17/28


H₂N-STAPPAHGVTSAPPDTRPAPGSTAPPK(Pal)G-OH



BP1-148

FIG. 17 Structure of lipopeptide **BP1-148**, a modified 25-amino-acid sequence derived from tumor-associated MUC1 mucin. BP1-148 is the antigen incorporated into the liposome formulation, together with an adjuvant, e.g. compound 1 - 6, to form the BLP25 liposomal vaccine.

18/28




R = CG, ATCGTT, GACGTT, ATCGAT, and any other ODN fragment.

Multi-valent effect is well known in many biological events. Very often the binding of a particular receptor to multi-ligands is the initial event of the multi-step activation cascade. The multi-arm of pentaerythritol (PET) provides an efficient scaffold to create a multi-valent system that will function as a more efficient ligand for a specific receptor. Cholesterol-conjugated to CpG containing ODN is expected to be potential ligands for TLR-9 and thus may have immunomodulatory activity.

FIG. 18 Clustered oligonucleotide - cholesterol conjugate

19/28

**FIG. 19** CpG-containing ODN lipidated at 3'-end

20/28





Fig. 20 Lipitated cyclic CpG dinucleotide analogues

21/28




Thymidine dimerization is one of the events behind UV activated DNA lesion. When this occurs, however, human body detects the damage and repair enzymes will be synthesized and the DNA damage will be repaired. It is likely that thymidine dimer may have functioned as a danger signal to which the immune system responds rapidly. Thus, those designed ODN analogues containing thymidine dimer, may have immune modulatory properties.

FIG. 21 Lipidated ODN containing thymidine dimer

22/28


Glycerol CpG



Unmethylated CpG dinucleotide with glycerol backbone: simple GNA derivatives designed as potential immune stimulatory agents.

FIG. 22 Lipitated glycerol CpG analogues


23/28



Lipo-teichoic acid (LTA) is the membrane component of Gram-positive bacteria. LTA is well known for its property to activate the innate immune response. Structures derived from LTA are therefore expected to have immune modulatory activities.

FIG. 23 Structures derived from lipo-teichoic acid

24/28



Modified Lipo-teichoic acid backbone, wherein the ester linkage between D-Alanine and the secondary hydroxyl group of glycerol unit is replaced by an amide bond, will be more stable toward hydrolytic condition.

FIG. 24 Modified LTA derivatives

25/28




FIG. 25 Hybrid structures derived from LTA, DNA, and GNA

26/28




FIG. 26

27/28

ds/sc Oligo(1:C)

Covalently linked oligo-inosinic acid and oligo-cytidylic acid have the potential to form single chain double strand structure that is expected to mimic the function of dsRNA as TLR-3 ligand.





FIG. 27

28/28

Two covalently linked oligo-inosinic -cytidinyl acid chains also have the potential to form double strand structure that may well mimic the function of dsRNA.



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.